



SureClean Plus

Column-Free Purification of DNA and dsRNA from PCR Reactions and Enzymatic Applications

Purification of nucleic acids is an essential technique for modern molecular biology. The removal of restriction enzymes, polymerases, dNTPs, primers and chemical inhibitors is an important factor for many downstream applications. As increasingly sensitive techniques are being developed, a higher demand is being placed on the ability to purify high-quality DNA. This application note reports on the performance of the Bioline reagent SureClean Plus, as a novel, column-free method for the purification of both DNA and dsRNA from PCR reactions or any enzymatic applications.




Figure 1. DNA and dsRNA purification using SureClean Plus. The Schematic diagram shows the simple, and cost-effective protocol used for purification of DNA using SureClean Plus.

REMOVAL OF PRIMERS FOLLOWING PCR REACTIONS

An important step in cloning PCR products is the removal of unincorporated dNTPs and primers, which are commonly over 25 bases in length, their presence can interfere with downstream applications and can have a detrimental effect on cloning experiments. To determine the cut-off point of fragment size by purification using SureClean Plus (fig. 1), we took a DNA ladder containing a range of sizes from 25 to 500bp, so as to compare SureClean Plus with two column-based purification methods. 30µl of HyperLadder 25bp (Cat. No. BIO-33031) was purified using the manufacturer's protocols. For each of the methods, DNA was resuspended in 30µl of TE, of which 5µl was loaded on to a 3.5% agarose gel. SureClean Plus efficiently purified DNA over 75bp (fig. 2). In contrast, the column-based method from competitor Q shows that following purification, DNA as small as 25bp was still present. Conversely, the method from competitor X efficiently removes primers, however, it also removes any genuine PCR products of under 200 bases. The data in fig. 2 shows that the level of recovery using competitors Q and X is lower than with SureClean Plus.

RECOVERY vs SIZE

SureClean Plus has been specifically developed to ensure the maximal recovery of nucleic acids after purification. The experiment below demonstrates the recovery efficiency on a wide range of different sizes of DNA. 30µl of HyperLadder 1kb (Cat No. BIO-33025) was purified using SureClean Plus and the column-based methods of Competitors Q and X. For each of the methods, DNA was resuspended in 30µl TE. 5µl was loaded on a 1% agarose gel (fig. 3).

These results clearly demonstrate the effectiveness of DNA recovery using SureClean Plus as compared with column-based purification methods.




Figure 2. DNA purification using SureClean Plus and Competitor's Q & X spin-column purification methods. 30µl of HyperLadder 25bp (Cat No. BIO-33031) was purified using the manufacturer's protocols. For each of the methods DNA was resuspended in 30µl TE, of which 5µl was loaded on to a 3.5% agarose gel.
Lane 1. HyperLadder 25bp
Lane 2. HyperLadder 25bp purified using SureClean Plus
Lane 3. HyperLadder 25bp purified using spin-columns from Competitor X
Lane 4. HyperLadder 25bp purified using spin-columns from Competitor Q




Figure 3. DNA purification using SureClean Plus and Competitor's Q & X spin-column purification methods. 30µl of HyperLadder 1kb (Cat No. BIO-33025) was purified using the manufacturer's protocols. For each of the methods DNA was resuspended in 30µl TE, of which 5µl was loaded on to a 1% agarose gel.
Lane 1. HyperLadder 1kb
Lane 2. HyperLadder 1kb purified using SureClean Plus
Lane 3. HyperLadder 1kb purified using spin-columns from Competitor X
Lane 4. HyperLadder 1kb purified using spin-columns from Competitor Q.



RECOVERY vs CONCENTRATION

In order to compare recovery rates with different DNA concentrations, solutions containing increasing amounts of DNA were purified using SureClean Plus and Competitor Q's spin-column purification (fig. 4). Unlike most column-based methods, SureClean Plus maximizes recovery with nucleic acid solutions of low, medium or high concentrations.

PURIFICATION OF dsRNA

In order to assess recovery of dsRNA, a 500bp fragment of dsRNA was synthesized using T7 RNA Transcription. This technique utilizes the synthesis of two single-stranded complementary RNA fragments, which are annealed to each other by heating to 65°C for 5 minutes and cooling slowly. Following the annealing step, the double-stranded fragment was purified using SureClean Plus in accordance with the standard purification protocol. The purified dsRNA was subsequently resuspended in DEPC-treated water (Cat No. BIO-38030), and analyzed on a 3.5% agarose (RNase-free) gel (fig. 5). This experiment demonstrates that SureClean Plus-purified the dsRNA with a good recovery rate.

REMOVAL OF RESTRICTION ENDONUCLEASES

Restriction enzyme digestion is a common method requiring downstream purification of DNA and the removal of enzyme. In order to test this, a fragment of the β -actin gene was amplified from human genomic DNA and then digested using *Pst*I enzyme (the β -actin fragment contained no *Pst*I sites). After a 2-hour incubation at 37°C, the gene was purified with SureClean Plus and with Competitor Q's column-based purification method. The purified fragment was then resuspended, following which λ DNA and *Pst*I buffer were added. Following a 2-hour incubation period at 37°C, no apparent digestion of the λ DNA was observed (fig. 6), so confirming the complete removal of the *Pst*I restriction enzyme.




Figure 4. Recovery vs concentration. DNA purification using SureClean Plus and Competitor's Q spin-column purification method. 30 μ l DNA solutions containing 100, 500, 1,000 and 2,500ng of plasmid DNA were purified using either SureClean Plus or Competitor Q spin-columns.




Figure 5. dsRNA recovery. A 500bp dsRNA fragment was synthesized using the T7 RNA Transcription kit, and then purified using SureClean Plus.

Lane 1. HyperLadder 25bp
Lane 2. Unpurified dsRNA
Lane 3. dsRNA purified using SureClean Plus




Figure 6. Enzyme Removal. An experiment was performed to remove the restriction endonucleases *Pst*I from a restriction digest and then demonstrate the absence of the enzyme in downstream applications.

Lane 1. HyperLadder 1kb
Lane 2. β -actin PCR fragment
Lane 3. β -actin PCR fragment plus *Pst*I and *Pst*I buffer
Lane 4. 20ng λ DNA
Lane 5. 20ng λ DNA plus *Pst*I and *Pst*I buffer
Lane 6. β -actin PCR fragment plus *Pst*I and *Pst*I buffer, purified using SureClean Plus. λ DNA and *Pst*I buffer were then added and incubated for 2 hours
Lane 7. β -actin PCR fragment plus *Pst*I and *Pst*I buffer, purified using Competitor Q's column based purification method. λ DNA and *Pst*I buffer were then added and incubated for 2 hours
Lane 8. HyperLadder 1kb



DOWNSTREAM APPLICATIONS


A. SEQUENCING

PCR products are used for numerous downstream applications, including sequencing. The length of read of a sequencing reaction depends on the purity of the DNA sample. In the following experiment, a 2Kb fragment was amplified by PCR and subsequently purified using SureClean Plus and Competitor U's single-tube purification method. The sequencing results obtained (fig. 7) indicate that SureClean Plus efficiently increased the length of read of the purified PCR product.

B. CLONING

The ability of SureClean Plus to enhance successful cloning was assessed. Three different-sized amplicons of 500bp, 1Kb and 2Kb DNA, were amplified using PCR from DNA. Following digestion of the PCR products with the restriction enzymes *EcoRI* and *HindIII*, the reaction was split into two halves. One half was further purified using SureClean Plus, whilst the other was left untreated. Each of the three amplicons was subsequently ligated into an *EcoRI/HindIII* plasmid vector (3:1 insert:vector ratio), using Quick-Stick Ligase (Cat No. BIO-27028). The ligation reaction was then directly cloned into α -Select Gold-Efficiency Competent Cells (Cat No. BIO-85027). The data shown (fig. 8) demonstrate the suitability of SureClean Plus as an important extra step in cloning the ligated PCR products.

SureClean Plus



Competitor U




Figure 7. Sequencing results of a PCR fragment purified by using SureClean Plus and Competitor U's single-tube purification method. Chart showing the length of read from a sequencing reaction, using SureClean Plus (S) and competitor U's single-tube method (U).

C. TRANSFECTION

A vector containing GFP was prepared with and without SureClean Plus as a final clean-up step after plasmid preparation. 3 μ g of the GFP vector was transiently transfected into CHOK1 cells and incubated for 24 hrs. Transfection efficiency was determined by flow cytometry on an Agilent 2100 Bioanalyzer. The data obtained (fig. 9) illustrate an increased transfection efficiency of the vector, following further purification of DNA using SureClean Plus.




Figure 8. Transformation efficiency of ligation reactions performed with unpurified and SureClean Plus-purified DNA. Three PCR amplicons, amplified from λ DNA were digested using the restriction endonucleases *EcoRI* and *HindIII*. The restriction digests were split into two. One half of the reaction was purified using SureClean Plus whereas the other half was not purified, and both the SureClean Plus-purified and the unpurified PCR fragments were ligated into an *EcoRI/HindIII* plasmid vector (3:1 insert:vector ratio) using Quick-Stick Ligase (Cat No. BIO-27028). The chart above shows an increase in colony-forming units (c.f.u) when the SureClean Plus-purified PCR amplicons for the ligation reaction are used.




Figure 9. Transfection efficiency of SureClean Plus-purified and unpurified GFP vector in CHOK1 cells. 3 μ g of a GFP vector, either purified with SureClean Plus after plasmid preparation or not purified was transiently transfected into CHOK1 cells and incubated for 24 hours. The transfection efficiency was determined by flow cytometry on an Agilent 2100 Bioanalyzer and the data are shown on a bar graph with the % of GFP-expressing cell population and % of CBNF-stained viable cells. MT is the mock transfection.



SUMMARY

Many of today's commercially available methods for purification of DNA rely on chaotropic salts and expensive spin-columns. These can lead to low recovery rates, DNA duplex denaturation and excessive cost for the researcher. In contrast, SureClean Plus provides column-free purification without organic solvents, resulting in high quality and purity of both DNA and dsRNA with negligible loss of product.

In addition, SureClean Plus is a universal purification system, efficiently removing primers/primer-dimers, restriction enzymes, unincorporated nucleotides and inhibitors of downstream applications, and conversely exhibiting almost no loss of product (recovery rates of up to 98%). SureClean Plus enables the researcher to resuspend the purified nucleic acid in any buffer and volume of choice, thus permitting the purification process to be tailored specifically to suit the experiment.

SureClean Plus enables standard 50µl reactions to be purified in a single tube at approximately one-third of the cost of competitors' spin-columns. When smaller reaction volumes are used, savings are even higher. For applications in which visual recognition of the DNA pellet is desired, SureClean Plus is supplied with a pink co-precipitant.

Please visit www.bioline.com to request a sample of SureClean Plus.

United Kingdom

Bioline Reagents Ltd
16 The Edge Business Centre
Humber Road
London NW2 6EW
United Kingdom
Tel: +44 (0)20 8830 5300
Fax: +44 (0)20 8452 2822

USA

Bioline USA Inc.
305 Constitution Dr.
Taunton, MA 02780
Toll Free: 888 257 5155
Tel: 508 880 8990
Fax: 508 880 8993

Germany

Bioline GmbH
Im Biotechnologiepark TGZ 2
D-14943 Luckenwalde
Tel: +49 (0)3371 681 229
Fax: +49 (0)3371 681 244

Australia

Bioline (Aust) Pty Ltd
PO Box 122
Alexandria NSW 1435
Australia
Tel: +61 (0)2 9209 4180
Fax: +61 (0)2 9209 4763